


AMENDMENTS TO THE CLAIMS

1. (Currently Amended) 4,4'-Bis(1,3,5-triazinylamino)stilbene-2,2'-disulfonic acid compound having the following formula:



in which

each of R^{11} and R^{12} independently is a hydrogen atom, an alkyl group having 1 to 20 carbon atoms, or an alkyl group having 1 to 20 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl, sulfo, and alkoxy hydrogen, methyl, ethyl, n-propyl, n-butyl, or 2-sulfoethyl;

R^{21} is a hydrogen atom, an alkyl group having 1 to 20 carbon atoms, an alkyl group having 1 to 20 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl, sulfo, and alkoxy, an aryl group having 6 to 20 carbon atoms, an aryl group having 6 to 20 carbon atoms which has one or more substituents selected from the group consisting of

~~hydroxyl, carboxyl, alkyl or alkoxy, or a group represented by the formula of L^1-CH_2OH wherein L^1 is an alkylene group having 2 to 8 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl and hydroxylalkyl having 1 to 3 carbon atoms or which has an intervening ether bonding,~~

~~R^{22} is a hydrogen atom, an alkyl group having 1 to 20 carbon atoms, an alkyl group having 1 to 20 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl, sulfo, and alkoxy, an aryl group having 6 to 20 carbon atoms, an aryl group having 6 to 20 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl, carboxyl, alkyl, or alkoxy, or a group represented by the formula of L^2CH_2OH wherein L^2 is an alkylene group having 2 to 8 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl and hydroxylalkyl having 1 to 3 carbon atoms or which has an intervening ether bonding; and~~






each of R^{21} and R^{22} independently is hydrogen, methyl, ethyl, n-propyl, isopropyl, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, 2,3-dihydroxypropyl, 2-sulfoethyl, 2-(2-hydroxyethoxy)ethyl, 2-[2-(2-hydroxyethoxy)ethoxy]ethyl, phenyl, naphthyl, 4-hydroxyphenyl, 3,5-dicarboxyphenyl, 4-methoxyphenyl, and 3-isopropylphenyl;

each of L^1 and L^2 is an alkylene group having 2 to 8 carbon atoms, which alkylene group has one or more substituents





selected from the group consisting of hydroxyl and hydroxyalkyl having 1 to 3 carbon atoms; and

M is a hydrogen atom, an alkali metal atom, ~~an alkaline earth metal atom, ammonium group,~~ or pyridinium group.

2. (Previously Presented) 4,4'-Bis(1,3,5-triazinylamino)stilbene-2,2'-disulfonic acid compound of claim 1, wherein at least one of L^1 and L^2 is a divalent group which is represented by one of the following formulas 1) to 5):



3. (Previously Presented) 4,4'-Bis(1,3,5-triazinylamino)stilbene-2,2'-disulfonic acid compound of claim 1, wherein at least one of L^1 and L^2 is a divalent group which is represented by one of the following formulas 1) to 4):




4. (Cancelled).

5. (Cancelled).

6. (Previously Presented) 4,4'-Bis(1,3,5-triazinylamino)stilbene-2,2'-disulfonic acid compound of claim 1, wherein each of R¹¹ and R¹² in the formula independently is a hydrogen or methyl.

7. (Previously Presented) 4,4'-Bis(1,3,5-triazinylamino)stilbene-2,2'-disulfonic acid compound of claim 1, wherein each of R²¹ and R²² in the formula independently is hydrogen, methyl, ethyl, isopropyl, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, 2,3-dihydroxypropyl, 2-(2-hydroxyethoxy)ethyl, 2-[2-(2-hydroxyethoxy)ethoxy]ethyl, phenyl, or 4-hydroxyphenyl.

8. (Currently Amended) An aqueous solution in which a 4,4'-Bis(1,3,5-triazinylamino)stilbene-2,2'-disulfonic acid compound having following formula is dissolved in water:



in which

each of R^{11} and R^{12} independently is ~~a hydrogen atom, an alkyl group having 1 to 20 carbon atoms, or an alkyl group having 1 to 20 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl, sulfo, and alkoxy~~ hydrogen, methyl, ethyl, n-propyl, n-butyl, or 2-sulfoethyl;

R^{21} ~~is a hydrogen atom, an alkyl group having 1 to 20 carbon atoms, an alkyl group having 1 to 20 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl, sulfo, and alkoxy, an aryl group having 6 to 20 carbon atoms, an aryl group having 6 to 20 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl, carboxyl, alkyl, or alkoxy, or a group represented by~~

~~the formula of L^1-CH_2OH wherein L^1 is an alkylene group having 2 to 8 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl and hydroxylalkyl having 1 to 3 carbon atoms or which has an intervening ether bonding;~~

~~R^{22} is a hydrogen atom, an alkyl group having 1 to 20 carbon atoms, an alkyl group having 1 to 20 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl, sulfo, and alkoxy, an aryl group having 6 to 20 carbon atoms, an aryl group having 6 to 20 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl, carboxyl, alkyl, or alkoxy, or a group represented by the formula of L^2-CH_2OH wherein L^2 is an alkylene group having 2 to 8 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl and hydroxylalkyl having 1 to 3 carbon atoms or which has an intervening ether bonding; and~~


each of R^{21} and R^{22} independently is hydrogen, methyl, ethyl, n-propyl, isopropyl, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, 2,3-dihydroxypropyl, 2-sulfoethyl, 2-(2-hydroxyethoxy)ethyl, 2-[2-(2-hydroxyethoxy)ethoxy]ethyl, phenyl, naphthyl, 4-hydroxyphenyl, 3,5-dicarboxyphenyl, 4-methoxyphenyl, and 3-isopropylphenyl;

each of L^1 and L^2 is an alkylene group having 2 to 8 carbon atoms, which alkylene group has one or more substituents

selected from the group consisting of hydroxyl and hydroxyalkyl having 1 to 3 carbon atoms; and

M is a hydrogen atom, an alkali metal atom, ~~an alkaline earth metal atom, ammonium group,~~ or pyridinium group.

9. (Currently Amended) A method of brightening a surface of material with fluorescence which comprises applying onto the surface an aqueous solution in which a 4,4'-Bis(1,3,5-triazinylamino)stilbene-2,2'-disulfonic acid compound having the following formula is dissolved in water:



in which

each of R^{11} and R^{12} independently is ~~a hydrogen atom, an alkyl group having 1 to 20 carbon atoms, or an alkyl group having 1 to 20 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl, sulfo, and alkoxy~~ hydrogen, methyl, ethyl, n-propyl, n-butyl, or 2-sulfoethyl;

~~R²¹ is a hydrogen atom, an alkyl group having 1 to 20 carbon atoms, an alkyl group having 1 to 20 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl, sulfo, and alkoxy, an aryl group having 6 to 20 carbon atoms, an aryl group having 6 to 20 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl, carboxyl, alkyl, or alkoxy, or a group represented by the formula of L¹-CH₂OH wherein L¹ is an alkylene group having 2 to 8 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl and hydroxylalkyl having 1 to 3 carbon atoms or which has an intervening ether bonding;~~

~~R²² is a hydrogen atom, an alkyl group having 1 to 20 carbon atoms, an alkyl group having 1 to 20 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl, sulfo, and alkoxy, an aryl group having 6 to 20 carbon atoms, an aryl group having 6 to 20 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl, carboxyl, alkyl, or alkoxy, or a group represented by the formula of L²-CH₂OH wherein L² is an alkylene group having 2 to 8 carbon atoms which has one or more substituents selected from the group consisting of hydroxyl and hydroxylalkyl having 1 to 3 carbon atoms or which has an intervening ether bonding; and~~

each of R²¹ and R²² independently is hydrogen, methyl, ethyl, n-propyl, isopropyl, 2-hydroxyethyl, 2-hydroxypropyl,

3-hydroxypropyl, 2,3-dihydroxypropyl, 2-sulfoethyl, 2-(2-hydroxyethoxy)ethyl, 2-[2-(2-hydroxyethoxy)ethoxy]ethyl, phenyl, naphthyl, 4-hydroxyphenyl, 3,5-dicarboxyphenyl, 4-methoxyphenyl, and 3-isopropylphenyl;

each of L¹ and L² is an alkylene group having 2 to 8 carbon atoms, which alkylene group has one or more substituents selected from the group consisting of hydroxyl and hydroxyalkyl having 1 to 3 carbon atoms; and

M is a hydrogen atom, an alkali metal atom, an alkaline earth metal atom, ammonium group, or pyridinium group.

10. (Cancelled).